Batty, G.D., Shipley, M.J., Marmot, M. & Davey Smith, G. Physical activity and cause-specific mortality in men with Type 2 diabetes/impaired glucose tolerance: evidence from the Whitehall study, 580
Bayly, G.R. *see* Lambert, A.P.
Beck-Nielsen, H. *see* Bentzen, J.
Bednarz, A.L. *see* Poulton, J.
Behrend, C. *see* Boehm, B.O.
Bellary, S. Exercise consultation in diabetes, 886
Beltramello, G.P. *see* Bruttomesso, D.
Bennetts, B.H. *see* Kao, Y.
Bentzen, J., Poulsen, P., Vaag, A., Beck-Nielsen, H. & Fenger, M. The influence of the polymorphism in apolipoprotein B codon 2488 on insulin and lipid levels in a Danish twin population, 12
Berger, D., Barroso, I., Soos, M., Yeo, G., Schafer, A.J., O'Rahilly, S. & Whitehead, J.P. Genetic variants of insulin receptor substrate-1 (IRS-1) in syndromes of severe insulin resistance. Functional analysis of Ala513Pro and Gly1158Glu IRS-1, 804
Berger, M. *see* Linkeschova, R.
Berger, M. *see* Neuvians, T.P.
Berger, M. *see* Trautner, C.
Berliner, S. *see* Elishkevitz, K.
Besser, R. 'Paint me a Hypo', 343
Betteridge, D.J. *see* Colhoun, H.M.
Bhattacharyya, A., Christodoulides, C., Kaushal, K., New, J.P. & Young, R.J. In-patient management of diabetes mellitus and patient satisfaction, 412
Bingley, P.J. *see* Douek, I.F.
Blackledge, H. *see* Hayward, L.M.
Blakytny, R. *see* Jude, E.B.
Bo, S., Cavallo-Perin, P., Gentile, L., Repetti, E. & Pagano, G. Type 2 diabetes with onset in the elderly is not a benign disease, 792
Bodansky, H.J. *see* Feltbower, R.G.
Boehm, B.O., Home, P.D., Behrend, C., Kamp, N.M. & Lindholm, A. Premixed insulin aspart 30 vs. premixed human insulin 30/70 twice daily: a randomized trial in Type 1 and Type 2 diabetic patients, 393
Bogardus, C. *see* Lindsay, R.S.
Bokemark, L., Wikstrand, J., Wedel, H. & Fagerberg, B. Insulin, insulin propeptides and intima-media thickness in the carotid artery in 58-year-old clinically healthy men. The Atherosclerosis and Insulin Resistance study (AIR), 144
Bolli, G.B. *see* Herz, M.
Bonadonna, R.C. *see* Bonora, E.
Bonora, E., Targher, G., Alberiche, M., Formentini, G., Calcaterra, F., Lombardi, S., Marini, F., Poli, M., Zenari, L., Raffaelli, A., Perbellini, S., Zenere, M.B., Saggiani, F., Bonadonna, R.C. & Muggeo, M. Predictors of insulin sensitivity in Type 2 diabetes mellitus, 535
Borch-Johnsen, K. IGT and IFG. Time for revision? 707
Borgo, E. *see* Trento, M.
Borretta, G. *see* Procopio, M.
Bosman, D.R., Osborne, C.A., Marsden, J.T., Macdougall, I.C., Gardner, W.N. & Watkins, P.J. Erythropoietin response to hypoxia in patients with diabetic autonomic neuropathy and non-diabetic chronic renal failure, 65
Bosnyák, Zs. *see* Tabák, Á.Gy.
Botha, J.L. *see* Hayward, L.M.
Bott, S. *see* Heise, T.
Bott, U. *see* Linkeschova, R.
Bougnères, P.F. *see* Coutant, R.
Boulton, A.J.M. *see* Abbott, C.A.
Boulton, A.J.M. *see* Armstrong, D.G.
Boulton, A.J.M. *see* Chaturvedi, N.
Boulton, A.J.M. *see* Jude, E.B.
Boulton, A.J.M. *see* Oyibo, S.O.
Boulton, A.J.M. *see* Paisley, A.N.
Bowen, K. *see* Kenardy, J.
Boyce, J.F. *see* Sinthanayothin, C.
Boyle, D.I.R. *see* Donnan, P.T.
Brailsford, S. *see* Davies, M.
Braspenning, J.C.C. *see* Frijling, B.D.
Bril, V. & Perkins, B.A. Comparison of vibration perception thresholds obtained with the Neurothesiometer and the CASE IV and relationship to nerve conduction studies, 661
Broadbent, D.M. *see* Younis, N.
Broers, S., le Cessie, S., van Vliet, K.P., Spinhoven, Ph., van der Ven, N.C.W. & Radder, J.K. Blood Glucose Awareness Training in Dutch Type 1 diabetes patients. Short-term evaluation of individual and group training, 157
Broomhall, J. *see* Donnan, P.T.
Brown, R.S. *see* Kipps, S.
Bruttomesso, D., Pianta, A., Crazzolara, D., Scaldaferri, E., Lora, L., Guarneri, G., Mongillo, A., Gennaro, R., Miola, M., Moretti, M., Confortin, L., Beltramello, G.P., Pais, M., Baritussio, A., Casiglia, E. & Tiengo, A. Continuous subcutaneous insulin infusion (CSII) in the Veneto region: efficacy, acceptability and quality of life, 628
Bruzzo, F. *see* Taverna, M.J.
Buch, H.N., Baskar, V., Barton, D.M., Kamalakannan, D., Akarca, C. & Singh, B.M. Combination of insulin and thiazolidinedione therapy in massively obese patients with Type 2 diabetes, 572
Bulmer, J. *see* Jude, E.B.
Burden, A.C. *see* Hayward, L.M.
Burden, M.L. *see* Hayward, L.M.
Butler, D. *see* Anderson, B.J.
Buysschaert, M. *see* Hermans, M.P.
Byrne, K. *see* McFarlane, S.I.

Calcaterra, F. *see* Bonora, E.
Calò, L. *see* Iori, E.
Canning, C. *see* Davies, M.
Cantero-Aguilar, L. *see* Coutant, R.
Carel, J.C. *see* Coutant, R.
Carr, M.E., Krishnaswami, A. & Martin, E.J. Platelet contractile force (PCF) and clot elastic modulus (CEM) are elevated in diabetic patients with chest pain, 862
Carrington, A.L. *see* Abbott, C.A.
Casiglia, E. *see* Bruttomesso, D.
Casson, I.F. *see* Platt, M.J.
Cassone-Faldetta, M. *see* Fava, D.
Cavallo, F. *see* Trento, M.
Cavallo-Perin, P. *see* Bo, S.
Cavan, D.A. *see* Kerr, D.
Ceolotto, G. *see* Iori, E.
Cesario, F. *see* Procopio, M.
le Cessie, S. *see* Broers, S.
Chan, A. *see* Kao, Y.
Chan, A.K.F. *see* Duffin, A.C.

Chan, J.C.N. *see* Ko, G.T.C.
Chan, J.C.N. *see* So, W.Y.
Chang, Y.F. *see* Hayward, L.M.
Chase, H.P., Saib, S.Z., MacKenzie, T., Hansen, M.M. & Garg, S.K. Post-prandial glucose excursions following four methods of bolus insulin administration in subjects with Type 1 diabetes, 317
Chaturvedi, N. Differing aspects of insulin resistance in diabetes complications: the shape of things to come. RD Lawrence lecture 2000, 973
Chaturvedi, N. *see* Kempler, P.
Chaturvedi, N., Abbott, C.A., Whalley, A., Widdows, P., Leggetter, S.Y. & Boulton, A.J.M. Risk of diabetes-related amputation in South Asians vs. Europeans in the UK, 99
Chevalier, A. *see* Taverna, M.J.
Chiarelli, F. *see* S. Mohn, A.
Cho, Y.M. *see* Kim, J.H.
Choi, D.S. *see* Choi, K.M.
Choi, K.M., Lee, J., Kim, D.R., Kim, S.K., Shin, D.H., Kim, N.H., Park, I.B., Choi, D.S. & Baik, S.H. Comparison of ADA and WHO criteria for the diagnosis of diabetes in elderly Koreans, 853
Chow, C.C. *see* So, W.Y.
Chowdury, T. & Lasker, S.S. Predicting CHD risk in patients with diabetes, 83
Christidis, D. *see* Liberopoulos, E.
Christodoulides, C. *see* Bhattacharyya, A.
Chuang, L.-M., Tsai, S.T., Huang, B.Y., Tai, T.Y. The status of diabetes control in Asia—a cross-sectional survey of 24 317 patients with diabetes mellitus in 1998, 978
Chung, S.S.M. *see* Liu, Y.F.
Close, C.F. *see* Lambert, A.P.
Clover, A. *see* Litwin, A.S.
Cockram, C.S. *see* So, W.Y.
Coker, R. *see* Shapo, L.
Colhoun, H.M., Thomason, M.J., Mackness, M.I., Maton, S.M., Betteridge, D.J., Durrington, P.N., Hitman, G.A., Neil, H.A.W. & Fuller, J.H. Design of the Collaborative AtoRvastatin Diabetes Study (CARDS) in patients with Type 2 diabetes, 201
Confortin, L. *see* Bruttomesso, D.
Connell, A. *see* Anderson, B.J.
Cook, H.L. *see* Sinthanayothin, C.
Coutant, R., Palmer, P., Carel, J.C., Cantero-Aguilar, L., Lebon, P., Bougnères, P.F. Detection of enterovirus RNA sequences in serum samples from autoantibody-positive subjects at risk for diabetes, 968
Crazzolara, D. *see* Bruttomesso, D.
Cundy, T. *see* Farrell, T.
Cundy, T. *see* Petrie, K.J.
Cundy, T., Slee, F., Gamble, G. & Neale, L. Hypertensive disorders of pregnancy in women with Type 1 and Type 2 diabetes, 482

D'Agostino Jr, R. *see* Festa, A.
Danne, T. *see* Kordonouri, O.
Daures, J.-P. *see* Richard, J.-L.
Davey Smith, G. *see* Batty, G.D.
Davies, M., Roderick, P., Canning, C. & Brailsford, S. The evaluation of screening policies for diabetic retinopathy using simulation, 762
Davis, G. *see* Ezenwaka, C.E.
Davis, T.M.E. *see* Balme, M.
Davis, W.A. *see* Balme, M.
Day, A.P. *see* Lambert, A.P.
Dayan, C.M. *see* Lambert, A.P.
De Luca, O. *see* Fava, D.
De Mattia, G. *see* Fava, D.
De Pergola, G. *see* Pannacciulli, N.
Deepa, R. *see* Mohan, V.
Deepa, R. *see* Rema, M.
Deepak, D., Ingham, D., Reynolds, R.M. & Walker, J.D. A case of euglycaemic diabetic ketoacidosis in pregnancy—a reply, 699
Deiss, D. *see* Kordonouri, O.
Dekker, J.M. *see* Adriaanse, M.C.
Dela, F. *see* Koning, E.J.P.
Delgado, J.A. *see* Abad, F.
Demaine, A.G. *see* Zhao, H.X.
Derfler, K. *see* Maca, T.H.
DeVries, J.H., Eskes, S.A., Snoek, F.J., Pouwer, F., van Ballegooie, E., Spijker, A.J., Kostense, P.J., Seubert, M. & Heine, R.J. Continuous intraperitoneal insulin infusion in patients with 'brittle' diabetes: favourable effects on glycaemic control and hospital stay, 496
Di Luzio, R. *see* S. Mohn, A.
Di Michele, *see* S. Mohn, A.
Díaz-Gómez, N.M. *see* Abad, F.
Didjurgeit, U., Kruse, J., Schmitz, N., Stückenschneider, P. & Sawicki, P.T. A time-limited, problem-orientated psychotherapeutic intervention in Type 1 diabetic patients with complications: a randomized controlled trial, 814
Dole, G. *see* Sampson, M.J.
Donaghue, K.C. *see* Duffin, A.C.
Donaghue, K.C. *see* Kao, Y.
Donnan, P.T. *see* Evans, J.M.M.
Donnan, P.T., Boyle, D.I.R., Broomhall, J., Hunter, K., MacDonald, T.M., Newton, R.W. & Morris, A.D. Prognosis following first acute myocardial infarction in Type 2 diabetes: a comparative population study, 448
Donnan, P.T., MacDonald, T.M. & Morris, A.D. Adherence to prescribed oral hypoglycaemic medication in a population of patients with Type 2 diabetes: a retrospective cohort study, 279
Donnan, P.T., Steinke, D.T., Newton, R.W. & Morris, A.D. Changes in treatment after the start of oral hypoglycaemic therapy in Type 2 diabetes: a population-based study, 606
Donovan, J. *see* Valabhji, J.
Dorchy, H. Improving outcomes for young people with diabetes, 702
Dorow, A. *see* Kordonouri, O.
Douek, I.F., Leech, N.J., Bingley, P.J. & Gale, E.A.M. Eczema and Type 1 diabetes, 174
Dressler, A. *see* Heise, T.
Duffin, A.C., Lam, A., Kidd, R., Chan, A.K.F., Donaghue, K.C. Ultrasonography of plantar soft tissues thickness in young people with diabetes, 1009
Duke, T. *see* Hayward, L.M.
Dunger, D.B. *see* Kipps, S.
Durrington, P.N. *see* Colhoun, H.M.
Durston, J. *see* Gill, G.
Dzien, A. *see* Lechleitner, M.
Dzien-Bischinger, C. *see* Lechleitner, M.

Eaton, S. *see* Kempler, P.

Edge, J.A. *see* Kipps, S.
Eguchi, K. *see* Takino, H.
Ehlin, A.G.C. *see* Montgomery, S.M.
Ehringer, H. *see* Maca, T.H.
van Eijk, J.T.M. *see* van de Poll-Franse, L.V.
Eisma, W.H. *see* Meijer, J.W.G.
Ekbom, A. *see* Montgomery, S.M.
El-Gammal, A. *see* Shaltout, A.A.
Eliasson, M., Lindahl, B., Lundberg, V. & Stegmayr, B. No increase in the prevalence of known diabetes between 1986 and 1999 in subjects 25–64 years of age in northern Sweden, 874
Elisaf, M. *see* Liberopoulos, E.
Elishkevitz, K., Fusman, R., Koffler, M., Shapira, I., Zeltser, D., Avitzour, D., Arber, N., Berliner, S. & Rotstein, R. Rheological determinants of red blood cell aggregation in diabetic patients in relation to their metabolic control, 152
Elkeles, R.S. *see* Valabhji, J.
Ellard, S. *see* Owen, K.R.
Ellingford, A. *see* Leese, G.P.
Ellis, J.D. *see* Leese, G.P.
Engström, G. & Janzon, L. Risk of developing diabetes in inversely related to lung function: a population-based cohort study, 167
Engström, G. *see* Hedblad, B.
Eskes, S.A. *see* DeVries, J.H.
Eskildsen, P.C. *see* Ihlemann, N.
Evans, J.M.M., Donnan, P.T. & Morris, A.D. Adherence to oral hypoglycaemic agents prior to insulin therapy in Type 2 diabetes, 685
Every, L.C. *see* Abbott, C.A.
Ezenwaka, C.E., Offiah, N.V., Kalloo, R. & Davis, G. The importance of diabetes health education in developing countries, 432

Fagerberg, B. *see* Bokemark, L.
Farrell, T., Neale, L. & Cundy, T. Congenital anomalies in the offspring of women with Type 1, Type 2 and gestational diabetes, 322
Fava, D., Cassone-Faldetta, M., Laurenti, O., De Luca, O., Ghiselli, A. & De Mattia, G. Gliclazide improves anti-oxidant status and nitric oxide-mediated vasodilation in Type 2 diabetes, 752
van der Feen, C., Neijens, F.S., Kanters, S.D.J.M., Mali, W.P.Th.M., Stolk, R.P. & Banga, J.D. Angiographic distribution of lower extremity atherosclerosis in patients with and without diabetes, 366
Feinle, C. *see* Horowitz, M
Feltbower, R.G., Bodansky, H.J., McKinney, P.A., Houghton, J., Stephenson, C.R. & Haigh, D. Trends in the incidence of childhood diabetes in south Asians and other children in Bradford, UK, 162
Felton, A. *see* Munro, N.
Fenger, M. *see* Bentzen, J.
Ferguson, M.W.J. *see* Jude, E.B.
Ferguson, S.C. *see* Herz, M.
Ferrannini, E. & Balkau, B. Insulin: in search of a syndrome, 724
Festa, A., D'Agostino Jr, R., Tracy, R.P. & Haffner, S.M. C-reactive protein is more strongly related to post-glucose load glucose than to fasting glucose in non-diabetic subjects; the Insulin Resistance Atherosclerosis Study, 939
Finnström, K. *see* Kullberg, C.E.
Fisher, M. Diabetes and Cardiovascular Disease, 260
Fisher, N.D.L. *see* Price, D.A.
Fitzpatrick, R. *see* Garratt, A.M.
Formentini, G. *see* Bonora, E.
Fowelin, J.H.R. *see* Johansson, K.E.A.
Fox, C.T. *see* Kipps, S.
François, L. *see* Martin, D.
Freeman, M.S., Mansfield, M.W., Barrett, J.H., Grant, P.J. Heritability of features of the insulin resistance syndrome in a community-based study of healthy families, 994
Frier, B.M. *see* Herz, M.
Frijling, B.D., Lobo, C.M., Hulscher, M.E.J.L., Akkermans, R.P., Braspenning, J.C.C., Prins, A., van der Wouden, J.C. & Grol, R.P.T.M. Multifaceted support to improve clinical decision making in diabetes care: a randomized controlled trial in general practice, 836
Fujiwara, Y. *see* Aso, Y.
Fukushima, M. *see* Nakai, Y.
Fuller, J.H. *see* Colhoun, H.M.
Fuller, J.H. *see* Kempler, P.
Fusman, R. *see* Elishkevitz, K.

Gale, E.A.M. *see* Douek, I.F.
Gamble, G. *see* Cundy, T.
Gandjour, A., Kleinschmit, F. & Lauterbach, K.W. European comparison of costs and quality in the prevention of secondary complications in Type 2 diabetes mellitus (2000–2001), 594
Gardner, W.N. *see* Bosman, D.R.
Garg, A. *see* Haque, W.A.
Garg, S.K. *see* Chase, H.P.
Garratt, A.M., Schmidt, L. & Fitzpatrick, R. Patient-assessed health outcome measures for diabetes: a structured review, 1
Gautier, J.F. *see* Sobngwi, E.
Ge, Y., MacDonald, D., Hait, H., Lipsky, B., Zasloff, M. & Holroyd, K. Microbiological profile of infected diabetic foot ulcers, 1032
Gennaro, R. *see* Bruttomesso, D.
Gentile, L. *see* Bo, S.
Gerel, B. *see* Suvd, J.
Ghiselli, A. *see* Fava, D.
Giani, G. *see* Trautner, C.
Gibbins, R. *see* Trento, M.
Gill-Randall, R. *see* Alcolado, J.C.
Gill, G., Durston, J., Johnson, R., MacLeod, K. & Watkins, P. Insulin-treated diabetes and driving in the UK, 435
Gill, G. Poor glycaemic control in female Type 1 adolescents—the 'fragile female diabetic' revisited? 699
Gill, G. The Diabetes Annual 13, 342
Gill, G.V. *see* Abbas, Z.G.
Gill, G.V. *see* Rotchford, A.P.
Giorgino, F. *see* Pannacciulli, N.
Giorgino, R. *see* Pannacciulli, N.
Gissler, M. *see* Vääräsmäki, M.
Goebel-Fabbri, A. *see* Anderson, B.J.
Gokal, R. *see* Oyibo, S.O.
Gough, K. *see* Strachan, M.W.J.
Grant, P.J. *see* Freeman, M.S.
Grasset, E. *see* Martin, D.
Green, B. *see* Kenardy, J.
Greene, S.A. *see* Howells, L.
Greenwood, R.H. *see* Sampson, M.J.
Gregory, R. *see* Swift, B.

Grenèche, M.O. *see* Martin, D.
Griffin, N.K. *see* Kipps, S.
Griffiths, J. *see* Abbott, C.A.
Grol, R.P.T.M. *see* Frijling, B.D.
Groothoff, J.W. *see* Meijer, J.W.G.
Grüters-Kieslich, A. *see* Kordonouri, O.
Gschwandtner, M.E. *see* Maca, T.H.
Guarneri, G. *see* Bruttomesso, D.
Gulam-Abbas, Z.G., Lutale, J.K., Morbach, S. & Archibald, L.K. Clinical outcome of diabetes patients hospitalized with foot ulcers, Dar es Salaam, Tanzania, 575
Gustafsson, J. *see* Tylleskär, K.
Gutniak, M.K. *see* Ahrén, B.
Guyot-Argenton, C. *see* Taverna, M.J.

Haastert, B. *see* Trautner, C.
Hadithi, M., Toorians, A. & Heine, R. The Houssay phenomenon, another clue to Sheehan's syndrome in a diabetic, 795
Haffner, S.M. *see* Festa, A.
Haigh, D. *see* Feltbower, R.G.
Hait, H. *see* Ge, Y.
Hamada, Y. *see* Takeuchi, N.
Hamilton, P. *see* Kerr, D.
Hampson, S.E. *see* Asimakopoulou, K.G.
Handley, G. *see* Wahid, S.T.
Hann, A.W. *see* Abbott, C.A.
Hanna, F.W.F. & Peters, J.R. Screening for gestational diabetes; past, present and future, 351
Hansen, M.M. *see* Chase, H.P.
Hanson, R.L. *see* Lindsay, R.S.
Haque, W.A., Vuitch, F., Garg, A. Post-mortem findings in familial partial lipodystrophy, Dunnigan variety, 1022
Hara, T. *see* Takeuchi, N.
Harding, S.P. *see* Younis, N.
Hardy, K.J. *see* Hulme, S.A.
Hartikainen, A.-L. *see* Vääräsmäki, M.
Harvey, I. *see* Sampson, M.J.
Hasegawa, G. *see* Kamiuchi, K.
Hasegawa, G. *see* Kitamura, A.
Hashizume, K. *see* Sato, Y.
Hattersley, A.T. *see* Owen, K.R.
Hawkins, P.N. *see* Swift, B.
Hayward, L.M., Burden, M.L., Burden, A.C., Blackledge, H., Raymond, N.T., Botha, J.L., Karwatowski, W.S.S., Duke, T. & Chang, Y.F. What is the prevalence of visual impairment in the general and diabetic populations: are there ethnic and gender differences? 27
Hedblad, B., Nilsson, P., Engström, G. & Janzon, L. Insulin resistance in non-diabetic subjects is associated with increased incidence of myocardial infarction and death, 470
Heine, R. *see* Hadithi, M.
Heine, R.J. *see* Adriaanse, M.C.
Heine, R.J. *see* DeVries, J.H.
Heine, R.J. *see* van de Poll-Franse, L.V.
Heinemann, L. *see* Heise, T.
Heise, T., Bott, S., Rave, K., Dressler, A., Rosskamp, R. & Heinemann, L. No evidence for accumulation of insulin glargine (LANTUS®): a multiple injection study in patients with Type 1 diabetes, 490
Heller, S. Mistaken priorities in diabetes research, 263
Hennessy, A.-R., Reynolds, R.M. & Walker, J.D. Blood glucose measurement in acute medicine: inadequate detection and management, 698
Hennessy, A.R. & Walker, J.D. Silent hypoglycaemia at the diabetic clinic, 261
Hermans, M.P., Van Leberghe, S. & Buysschaert, M. The [C282Y/wt] heterozygous HFE genotype is associated with lower blood pressure and HDL-cholesterol in Type 2 diabetes, 796
Hernández, C., Lecube, A., Segura, R.M., Sararols, L. & Simó, R. Nitric oxide and vascular endothelial growth factor concentrations are increased but not related in vitreous fluid of patients with proliferative diabetic retinopathy, 655
Herold, M. *see* Lechleitner, M.
Herz, M., Arora, V., Sun, B., Ferguson, S.C., Bolli, G.B. & Frier, B.M. Basal-bolus insulin therapy in Type 1 diabetes: comparative study of pre-meal administration of a fixed mixture of insulin lispro (50%) and neutral protamine lispro (50%) with human soluble insulin, 917
Heyburn, P.J. *see* Sampson, M.J.
Hilsted, J. *see* Rosenfalck, A.M.
Hiramatsu, K. *see* Sato, Y.
Hitman, G.A. *see* Colhoun, H.M.
Ho, L.T. *see* Kuo, C.S.
Hodgkins, P.R. *see* Litwin, A.S.
Hodson, M.E. *see* Yung, B.
Hollenberg, N.K. *see* Price, D.A.
Holroyd, K. *see* Ge, Y.
Holst, J.J. *see* Ahrén, B.
Holt, T.A. A chaotic model for tight diabetes control, 274
Home, P.D. *see* Boehm, B.O.
Home, P.D. *see* Marshall, S.M.
Hooper, J. *see* Yung, B.
Hoppichler, F. *see* Lechleitner, M.
Hörl, W.H. *see* Maca, T.H.
Horowitz, M., O'Donovan, D., Jones, K.L., Feinle, C., Rayner, C.K. & Samson, M. Gastric emptying in diabetes: clinical significance and treatment, 177
Hotta, N. *see* Kondo, K.
Hotta, N. *see* Takeuchi, N.
Houghton, J. *see* Feltbower, R.G.
Howard, C.V. *see* Platt, M.J.
Howells, L., Wilson, A.C., Skinner, T.C., Newton, R., Morris, A.D. & Greene, S.A. A randomized control trial of the effect of negotiated telephone support on glycaemic control in young people with Type 1 diabetes, 643
Howlett, H. *see* Marre, M.
Hsiao, L.C. *see* Kuo, C.S.
Huang, B.Y. *see* Chuang, L.-M.
Huh, K.B. *see* Ahn, C.W.
Hulme, S.A., Tin-U, A., Hardy, K.J. & Joyce, P.W. Evaluation of a district-wide screening programme for diabetic retinopathy utilizing trained optometrists using slit-lamp and Volk lenses, 741
Hulscher, M.E.J.L. *see* Frijling, B.D.
Humphries, J. *see* Sampson, M.J.
Hunt, M.A. *see* Lambert, A.P.
Hunter, K. *see* Donnan, P.T.
Hussein, A. *see* Abbott, C.A.
Hwu, C.M. *see* Kuo, C.S.

Ichikawa, K. *see* Sato, Y.
Ihlemann, N., Stokholm, K.H. & Eskildsen, P.C. Impaired vascular reactivity is present despite normal levels of von Willebrand factor in patients with uncomplicated Type 2 diabetes, 476
Inagaki, A. *see* Kato, Y.
Ingham, D. *see* Deepak, D.
Inukai, G. *see* Matsumoto, K.
Inukai, T. *see* Aso, Y.
Iori, E., Calò, L., Valbusa, D., Ceolotto, G., Milani, M., Pengo, V., de Kreutzenberg, S.V., Tiengo, A. & Avogaro, A. Diabetic ketosis activates lymphomonocyte-inducible nitric oxide synthase, 777
Ishibashi, S. *see* Nagasaka, S.
Ishii, M. *see* Kamiuchi, K.
Ishii, M. *see* Kitamura, A.
Ishikawa, S. *see* Nagasaka, S.
Itatsu, T. *see* Kato, Y.

Jackson, N. *see* Abbott, C.A.
Jackson, N.J. *see* Oyibo, S.O.
James, E. *see* Oyibo, S.O.
Jansen van Rosendaal, A. *see* Abbink, E.J.
Janzon, L. *see* Engström, G.
Janzon, L. *see* Hedblad, B.
Jeon, H.J. *see* Kim, J.H.
Johansson, K.E.A., Marklund, B.R.G. & Fowelin, J.H.R. Evaluation of a new screening method for detecting peripheral arterial disease in a primary health care population of patients with diabetes mellitus, 307
John, W.G. *see* Marshall, S.M.
Johnson, K.E. *see* Abbott, C.A.
Johnson, R. *see* Gill, G.
Joner, G. *see* Stene, L.C.
Jones, K.L. *see* Horowitz, M
Jones, S.D. *see* Manning, P.J.
de Jong, S.A. *see* Manning, P.J.
Jousilahti, P. *see* Al-Lawati, J.A.
Joyce, P.W. *see* Hulme, S.A.
Jude, E.B. *see* Oyibo, S.O.
Jude, E.B., Blakytny, R., Bulmer, J., Boulton, A.J.M. & Ferguson, M.W.J. Transforming growth factor-beta 1, 2, 3 and receptor type I and II in diabetic foot ulcers, 440

Kalloo, R. *see* Ezenwaka, C.E.
Kamalakannan, D. *see* Buch, H.N.
Kamiuchi, K. *see* Kitamura, A.
Kamiuchi, K., Hasegawa, G., Obayashi, H., Kitamura, A., Ishii, M., Yano, M., Kanatsuna, T., Yoshikawa, T. & Nakamura, N. Intercellular adhesion molecule-1 (ICAM-1) polymorphism is associated with diabetic retinopathy in Type 2 diabetes mellitus, 371
Kamp, N.M. *see* Boehm, B.O.
Kanai, A. *see* Takeuchi, N.
Kanatsuna, T. *see* Kamiuchi, K.
Kanazawa, Y. *see* Takino, H.
Kang, B.S. *see* Kim, J.H.
Kanters, S.D.J.M. *see* van der Feen, C.
Kao, Y., Donaghue, K.C., Chan, A., Bennetts, B.H., Knight, J. & Silink, M. Paraoxonase gene cluster is a genetic marker for early microvascular complications in Type 1 diabetes, 212
Karvonen, M. *see* Shaltout, A.A.
Karwatowski, W.S.S. *see* Hayward, L.M.
Katakura, M. *see* Sato, Y.
Kato, Y., Miura, Y., Inagaki, A., Itatsu, T. & Oiso, Y. Age of onset posssibly associated with the degree of heteroplasmy in two male siblings with diabetes mellitus having an A to G transition at 3243 of mitochondrial DNA, 784
Katzenschlager, R. *see* Maca, T.H.
Kaushal, K. *see* Bhattacharyya, A.
Kawakami, A. *see* Nagasaka, S.
Kawamura, T. *see* Takeuchi, N.
Kawasaki, E. *see* Takino, H.
Kemp, M. *see* Yung, B.
Kempler, P., Tesfaye, S., Chaturvedi, N., Stevens, L.K., Webb, D.J., Eaton, S., Kerényi, Zs., Tamás, Gy., Ward, J.D. & Fuller, J.H. Autonomic neuropathy is associated with increased cardiovascular risk factors: the EURODIAB IDDM Complications Study, 900
Kenardy, J., Mensch, M., Bowen, K., Green, B. & Walton, J. Group therapy for binge eating in Type 2 diabetes: a randomized trial, 234
Kerényi, Zs. *see* Kempler, P.
Kerényi, Zs. *see* Tabák, Á.Gy.
Kerr, D., Hamilton, P., Cavan, D.A. Preventing glycaemic excursions in diabetic patients requiring percutaneous endoscopic gastrostomy (PEG) feeding after a stroke, 1006
Kerr, D., Nicholls, H. & Cavan, D.A. The lost tribes in diabetes, 703
Kidd, R. *see* Duffin, A.C.
Kim, D.R. *see* Choi, K.M.
Kim, H.S. *see* Ahn, C.W.
Kim, J.H., Park, K.S., Cho, Y.M., Kang, B.S., Kim, S.K., Jeon, H.J., Kim, S.Y. & Lee, H.K. The prevalence of the mitochondrial DNA 16189 variant in non-diabetic Korean adults and its association with higher fasting glucose and body mass index, 681
Kim, K.R. *see* Ahn, C.W.
Kim, N.H. *see* Choi, K.M.
Kim, S.K. *see* Choi, K.M.
Kim, S.K. *see* Kim, J.H.
King, H. *see* Ramachandran, A.
King, H. *see* Suvd, J.
Kipps, S., Bahu, T., Ong, K., Ackland, F.M., Brown, R.S., Fox, C.T., Griffin, N.K., Knight, A.H., Mann, N.P., Neil, H.A.W., Simpson, H., Edge, J.A. & Dunger, D.B. Current methods of transfer of young people with Type 1 diabetes to adult services, 649
Kishimoto, H. *see* Nakai, Y.
Kitamura, A. *see* Kamiuchi, K.
Kitamura, A., Hasegawa, G., Obayashi, H., Kamiuchi, K., Ishii, M., Yano, M., Tanaka, T., Yamaguchi, M., Shigeta, H., Ogata, M., Nakamura, N., Yoshikawa, T. Interleukin-6 polymorphism (−634C/G) in the promotor region and the progression of diabetic nephropathy in Type 2 diabetes, 1000
Kleinschmit, F. *see* Gandjour, A.
Knight, A.H. *see* Kipps, S.
Knight, J. *see* Kao, Y.
Knip, M. *see* Riihimaa, P.H.
Knowler, W.C. *see* Lindsay, R.S.
Ko, B.C.B. *see* Liu, Y.F.
Ko, G.T.C. *see* So, W.Y.
Ko, G.T.C., Tam, W.H., Chan, J.C.N. & Rogers, M. Prevalence of

gestational diabetes mellitus in Hong Kong based on the 1998 WHO criteria, 80
Koffler, M. *see* Elishkevitz, K.
Komatsu, M. *see* Sato, Y.
Kondo, K., Matsubara, T., Nakamura, J. & Hotta, N. Characteristic patterns of circadian variation in plasma catecholamine levels, blood pressure and heart rate variability in Type 2 diabetic patients, 359
Koning, E.J.P., Visboll, T. & Dela, F. 19th Anglo–Danish–Dutch Diabetes Group meeting, 259
Koopmanschap, M.A. *see* Redekop, W.K.
Koppensteiner, R. *see* Maca, T.H.
Kordonouri, O., Deiss, D., Danne, T., Dorow, A., Bassir, C. & Grüters-Kieslich, A. Predictivity of thyroid autoantibodies for the development of thyroid disorders in children and adolescents with Type 1 diabetes, 518
Koshima, I. *see* Usui, H.
Kostense, P.J. *see* DeVries, J.H.
de Kreutzenberg, S.V. *see* Iori, E.
Krishnaswami, A. *see* Carr, M.E.
Krolewski, A.S. *see* Weijnen, C.F.
Kroon, M. *see* Särnblad, S.
Kruse, J. *see* Didjurgeit, U.
Kullberg, C.E., Abrahamsson, M., Arnqvist, H.J., Finnström, K. & Ludvigsson, J. Prevalence of retinopathy differs with age at onset of diabetes in a population of patients with Type 1 diabetes, 924
Kuo, C.S., Hwu, C.M., Kwok, C.F., Hsiao, L.C., Weih, M.J., Lee, S.H., Lee, Y.S. & Ho, L.T. Surrogate estimates of insulin sensitivity in Chinese diabetic patients and their offspring, 735
Kusaka, I. *see* Nagasaka, S.
Kwok, C.F. *see* Kuo, C.S.
Kynde, K. *see* Ødum, L.

Laaksonen, D.E., Lakka, T.A., Lakka, H.-M., Nyyssönen, K., Rissanen, T., Niskanen, L.K. & Salonen, J.T. Serum fatty acid composition predicts development of impaired fasting glycaemia and diabetes in middle-aged men, 456
Laffel, L.M.B. *see* Anderson, B.J.
Laffel, L.M.B. *see* Price, D.A.
Laing, I. *see* Oyibo, S.O.
Laing, I. *see* Yates, A.P.
Laji, K. *see* Alcolado, J.C.
Lakka, H.-M. *see* Laaksonen, D.E.
Lakka, T.A. *see* Laaksonen, D.E.
Lal, S. *see* Sinthanayothin, C.
Lam, A. *see* Duffin, A.C.
Lam, K.S.L. *see* Liu, Y.F.
Lam, K.S.L. *see* Tan, K.C.B.
Lambert, A.P. & Close, C.F. Charcot neuroarthropathy of the knee in Type 1 diabetes: treatment with total knee arthroplasty, 338
Lambert, A.P., Hunt, M.A., Day, A.P., Bayly, G.R. & Dayan, C.M. Reproducibility of individualized coronary heart disease risk calculations in patients with diabetes mellitus, 514
Lansang, M.C. *see* Price, D.A.
Lant, A.F *see* Yung, B.
Lasker, S.S. *see* Chowdury, T.
Laurenti, O. *see* Fava, D.
Lauterbach, K.W. *see* Gandjour, A.
Leblanc, H. *see* Sobngwi, E.
Lebon, P. *see* Coutant, R.
Lechleitner, M., Herold, M., Dzien-Bischinger, C., Hoppichler, F., Dzien, A. Tumour necrosis factor-alpha plasma levels in elderly patients with Type 2 diabetes mellitus—observations over 2 years, 949
Lecube, A. *see* Hernández, C.
Lee, H.C. *see* Ahn, C.W.
Lee, H.K. *see* Kim, J.H.
Lee, J. *see* Choi, K.M.
Lee, S.H. *see* Kuo, C.S.
Lee, Y.S. *see* Kuo, C.S.
Leech, N.J. *see* Douek, I.F.
Leese, G.P., Ellis, J.D., Morris, A.D. & Ellingford, A. Does direct ophthalmoscopy improve retinal screening for diabetic eye disease by retinal photography? 867
Leggetter, S.Y. *see* Chaturvedi, N.
Legrand, C. *see* Martin, D.
Lehert, P. *see* Marre, M.
Leong, K.S. *see* Li Voon Chong, J.S.W.
Lepage, V. *see* Sobngwi, E.
Levi, Z. *see* Rachmani, R.
Levy, V. *see* Sobngwi, E.
Li, J.K.Y. *see* So, W.Y.
Li Voon Chong, J.S.W., Leong, K.S., Wallymahmed, M., Sturgess, R. & MacFarlane, I.A. Is coeliac disease more prevalent in young adults with coexisting Type 1 diabetes mellitus and autoimmune thyroid disease compared with those with Type 1 diabetes mellitus alone? 334
Liberopoulos, E., Christidis, D. & Elisaf, M. Marked hyperphosphataemia in a patient with diabetic ketoacidosis, 885
Licha-Müntz, G. *see* Martin, D.
Lim, S.K. *see* Ahn, C.W.
Lindahl, B. *see* Eliasson, M.
Lindblad, U. *see* Östgren, C.J.
Lindholm, A. *see* Boehm, B.O.
Lindow, S.W. *see* Rao, S.
Lindsay, R.S., Prochazka, M., Baier, L.J., Knowler, W.C., Bogardus, C. & Hanson, R.L. Currently identified genes affecting insulin resistance are not associated with birth weight in the Pima population, 882
Linkeschova, R., Raoul, M., Bott, U., Berger, M. & Spraul, M. Less severe hypoglycaemia, better metabolic control, and improved quality of life in Type 1 diabetes mellitus with continuous subcutaneous insulin infusion (CSII) therapy; an observational study of 100 consecutive patients followed for a mean of 2 years, 746
Links, T.P. *see* Meijer, J.W.G.
Lipsky, B. *see* Ge, Y.
Litwin, A.S., Clover, A., Hodgkins, P.R. & Luff, A.J. Affluence is not related to delay in diagnosis of Type 2 diabetes as judged by the development of diabetic retinopathy, 843
Liu, Y.F., Wat, N.M.S., Chung, S.S.M., Ko, B.C.B. & Lam, K.S.L. Diabetic nephropathy is associated with the 5′-end dinucleotide repeat polymorphism of the aldose reductase gene in Chinese subjects with Type 2 diabetes, 113
Lobo, C.M. *see* Frijling, B.D.
Lombardi, S. *see* Bonora, E.
Lora, L. *see* Bruttomesso, D.
Ludvigsson, J. *see* Kullberg, C.E.
Luff, A.J. *see* Litwin, A.S.
Lundberg, V. *see* Eliasson, M.

Lutale, J.K. *see* Gulam-Abbas, Z.G.
Lutterman, J.A. *see* Abbink, E.J.
Lyaruu, I. *see* Neuhann, H.F.

Maca, T.H., Ahmadi, R., Derfler, K., Ehringer, H., Gschwandtner, M.E., Hörl, W.H., Katzenschlager, R., Müller-Knespel, E., Koppensteiner, R., Schneider, B., Stümpflen, A., Ugurluoglu, A. & Minar, E. Influence of lipoprotein(a) on restenosis after femoropopliteal percutaneous transluminal angioplasty in Type 2 diabetic patients, 300
Macaulay, V.A. *see* Poulton, J.
MacDonald, D. *see* Ge, Y.
MacDonald, T.M. *see* Donnan, P.T.
Macdougall, I.C. *see* Bosman, D.R.
MacFarlane, I.A. *see* Li Voon Chong, J.S.W.
Machattie, T. *see* Rotchford, A.P.
MacKenzie, T. *see* Chase, H.P.
Mackness, M.I. *see* Colhoun, H.M.
MacLeod, K. *see* Gill, G.
Madarász, E. *see* Tabák, Á.Gy.
Madsbad, S. *see* Rosenfalck, A.M.
Magnus, P. *see* Stene, L.C.
Magro, G. *see* Procopio, M.
Makino, H. *see* Usui, H.
Mali, W.P.Th.M. *see* van der Feen, C.
Manley, S.E. *see* Marshall, S.M.
Mann, N.P. *see* Kipps, S.
Manning, P.J., Sutherland, W.H.F., Allum, A.R., de Jong, S.A. & Jones, S.D. Effect of hormone replacement therapy on inflammation-sensitive proteins in post-menopausal women with Type 2 diabetes, 847
Mansell, P. Diabetes and driving, 617
Mansfield, M.W. *see* Freeman, M.S.
Marini, F. *see* Bonora, E.
Marklund, B.R.G. *see* Johansson, K.E.A.
Marmot, M. *see* Batty, G.D.
Marre, M., Howlett, H., Lehert, P. & Allavoine, T. Improved glycaemic control with metformin–glibenclamide combined tablet therapy (Glucovance®) in Type 2 diabetic patients inadequately controlled on metformin, 673
Marsden, J.T. *see* Bosman, D.R.
Marshall, S.M., Home, P.D., Manley, S.E., Barth, J.H. & John, W.G. Standardization of glycated haemoglobin, 429
Martin, D., Licha-Müntz, G., Grasset, E., Grenèche, M.O., Nouet, D., François, L., Legrand, C., Polak, M., Augendre-Ferrante, B., Tubiana-Rufi, N., Robert, J.J. Efficacy of Humalog® injections before an afternoon meal and their acceptance by children and adolescents with Type 1 diabetes, 1026
Martin, E.J. *see* Carr, M.E.
Masson, E.A. *see* Rao, S.
Maton, S.M. *see* Colhoun, H.M.
Matsubara, T. *see* Kondo, K.
Matsuda, M. *see* Usui, H.
Matsumoto, K., Sera, Y., Ueki, Y., Inukai, G., Niiro, E. & Miyake, S. Comparison of serum concentrations of soluble adhesion molecules in diabetic microangiopathy and macroangiopathy, 822
Matthews, D.R. *see* Wallace, T.M.
Maurino, M. *see* Trento, M.
Mauvais-Jarvis, F. *see* Sobngwi, E.
Mbanya, J.C. *see* Sobngwi, E.
McAllister, I. *see* Balme, M.
McColl, A.J. *see* Valabhji, J.
McFarlane, S.I., Byrne, K., Shin, J. & Williams, R. Intussusception in an adult patient with severe hyperglycaemia—a case report, 611
McIntosh, C. *see* Munro, N.
McKee, M. *see* Shapo, L.
McKendrick, O. *see* Platt, M.J.
McKinney, P.A. *see* Feltbower, R.G.
McKnight, J.A. *see* Strachan, M.W.J.
Mclay, L. *see* Oyibo, S.O.
Meigs, J.B. *see* Weijnen, C.F.
Meijer, J.W.G., Smit, A.J., Sonderen, E.V., Groothoff, J.W., Eisma, W.H., Links, T.P. Symptom scoring systems to diagnose distal polyneuropathy in diabetes: the Diabetic Neuropathy Symptom score, 962
Melander, A. *see* Östgren, C.J.
Melchionda, N. *see* Serrano-Rios, M.
Mensah, E. *see* Sinthanayothin, C.
Mensch, M. *see* Kenardy, J.
Milani, M. *see* Iori, E.
Millward, B.A. *see* Zhao, H.X.
Minar, E. *see* Maca, T.H.
Miola, M. *see* Bruttomesso, D.
Miura, Y. *see* Kato, Y.
Miyake, S. *see* Matsumoto, K.
Mohammed, A.J. *see* Al-Lawati, J.A.
Mohan, V. *see* Rema, M.
Mohan, V. & Deepa, R. Author's reply, 81
Mohn, A., Di Michele, S., Di Luzio, R., Tumini, S. & Chiarelli, F. The effects of subclinical hypothyroidism on metabolic control in children and adolescents with Type 1 diabetes mellitus, 70
Molineri, N. *see* Procopio, M.
Mongillo, A. *see* Bruttomesso, D.
Montgomery, S.M., Ehlin, A.G.C., Ekbom, A., Wakefield, A.J. Pertussis infection in childhood and subsequent Type 1 diabetes mellitus, 986
Morbach, S. *see* Gulam-Abbas, Z.G.
Moreno-Carretero, E. *see* Serrano-Rios, M.
Moretti, M. *see* Bruttomesso, D.
Morris, A.D. *see* Donnan, P.T.
Morris, A.D. *see* Evans, J.M.M.
Morris, A.D. *see* Howells, L.
Morris, A.D. *see* Leese, G.P.
Morrish, N.J. *see* Asimakopoulou, K.G.
Moussa, M.A.A. *see* Shaltout, A.A.
Moyeed, R.A. *see* Zhao, H.X.
Msuya, L. *see* Neuhann, H.F.
Muggeo, M. *see* Bonora, E.
Müller-Knespel, E. *see* Maca, T.H.
Munro, N., Felton, A. & McIntosh, C. Is multidisciplinary learning effective among those caring for people with diabetes? 799

Nagasaka, S. *see* Nakai, Y.
Nagasaka, S., Abe, T., Kawakami, A., Kusaka, I., Nakamura, T., Ishikawa, S., Saito, T. & Ishibashi, S. Pioglitazone-induced hepatic injury in a patient previously receiving troglitazone with success, 347
Nagataki, S. *see* Takino, H.
Nagy, E. *see* Tabák, Á.Gy.

Nakai, Y., Fukushima, M., Nakaishi, S., Kishimoto, H., Seino, Y., Nagasaka, S., Sakai, M. & Taniguchi, A. The threshold value for insulin resistance on homeostasis model assessment of insulin sensitivity, 346
Nakaishi, S. *see* Nakai, Y.
Nakamura, J. *see* Kondo, K.
Nakamura, J. *see* Takeuchi, N.
Nakamura, N. *see* Kamiuchi, K.
Nakamura, N. *see* Kitamura, A.
Nakamura, N. *see* Takeuchi, N.
Nakamura, T. *see* Nagasaka, S.
Nam, J.H. *see* Ahn, C.W.
Neale, L. *see* Cundy, T.
Neale, L. *see* Farrell, T.
Neijens, F.S. *see* van der Feen, C.
Neil, H.A.W. *see* Colhoun, H.M.
Neil, H.A.W. *see* Kipps, S.
Neuhann, H.F., Warter-Neuhann, C., Lyaruu, I. & Msuya, L. Diabetes care in Kilimanjaro region: clinical presentation and problems of patients of the diabetes clinic at the regional referral hospital—an inventory before structured intervention, 509
Neuvians, T.P. & Berger, M. Diabetes care in cats and dogs, 77
New, J.P. *see* Bhattacharyya, A.
Newton, R. *see* Howells, L.
Newton, R.W. *see* Donnan, P.T.
Nicholls, A.J. The impact of atherosclerotic renovascular disease on diabetic renal failure, 889
Nicholls, H. *see* Kerr, D.
Niessen, L.W. *see* Redekop, W.K.
Niiro, E. *see* Matsumoto, K.
Nilsson, P. *see* Hedblad, B.
Niskanen, L.K. *see* Laaksonen, D.E.
Njølstad, P.R. *see* Sagen, J.V.
Noormohamed, F.H. *see* Yung, B.
Nouet, D. *see* Martin, D.
Nyyssönen, K. *see* Laaksonen, D.E.

O'Donovan, D. *see* Horowitz, M
O'Rahilly, S. *see* Berger, D.
Obayashi, H. *see* Kamiuchi, K.
Obayashi, H. *see* Kitamura, A.
Ødum, L. & Kynde, K. Handling of blood samples for diagnosis of diabetes mellitus, 345
Offiah, N.V. *see* Ezenwaka, C.E.
Ogata, M. *see* Kitamura, A.
Ohfusa, H. *see* Sato, Y.
Oishi, T. *see* Usui, H.
Oiso, Y. *see* Kato, Y.
Ong, K. *see* Kipps, S.
Osborne, C.A. *see* Bosman, D.R.
Osei, S.Y. *see* Price, D.A.
Östgren, C.J., Lindblad, U., Ranstam, J., Melander, A. & Råstam, L. Glycaemic control, disease duration and β-cell function in patients with Type 2 diabetes in a Swedish community. Skaraborg Hypertension and Diabetes Project, 125
Otgooloi, H. *see* Suvd, J.
Owen, K.R., Shepherd, M., Stride, A., Ellard, S. & Hattersley, A.T. Heterogeneity in young adult onset diabetes: aetiology alters clinical characteristics, 758
Owens, D.R. *see* Trento, M.
Oyibo, S.O., Prasad, Y.D.M., Jackson, N.J., Jude, E.B. & Boulton, A.J.M. The relationship between blood glucose excursions and painful diabetic peripheral neuropathy: a pilot study, 870
Oyibo, S.O., Pritchard, G.M., Mclay, L., James, E., Laing, I., Gokal, R. & Boulton, A.J.M. Blood glucose overestimation in diabetic patients on continuous ambulatory peritoneal dialysis for end-stage renal disease, 693

Pacher, N. *see* Taverna, M.J.
Padfield, P.L. *see* Strachan, M.W.J.
Pagano, G. *see* Bo, S.
Pais, M. *see* Bruttomesso, D.
Paisley, A.N., Abbott, C.A., van Schie, C.H.M. & Boulton, A.J.M. A comparison of the Neuropen against standard quantitative sensory-threshold measures for assessing peripheral nerve function, 400
Palmer, P. *see* Coutant, R.
Pandit, R.J. & Taylor, R. Quality assurance in screening for sight-threatening diabetic retinopathy, 285
Pang, R.W.C. *see* Tan, K.C.B.
Pannacciulli, N., De Pergola, G., Giorgino, F. & Giorgino, R. A family history of Type 2 diabetes is associated with increased plasma levels of C-reactive protein in non-smoking healthy adult women, 689
Parer-Richard, C. *see* Richard, J.-L.
Park, I.B. *see* Choi, K.M.
Park, K.S. *see* Kim, J.H.
Passera, P. *see* Trento, M.
Pengo, V. *see* Iori, E.
Pennycook, S. *see* Platt, M.J.
Perbellini, S. *see* Bonora, E.
Pérez, R. *see* Abad, F.
Perkins, B.A. *see* Bril, V.
Perlemuter, L. & Yomtov, B. Feasibility and usefulness of dedicated software and e-mail for self-monitoring blood glucose in treating diabetes, 701
Peters, J.R. *see* Hanna, F.W.F.
Petrella, E. *see* Ahrén, B.
Petrie, K.J. & Cundy, T. Alternative medicine recommendations to patients with Type 2 diabetes visiting health product shops and pharmacies, 1035
Pia, A. *see* Procopio, M.
Pianta, A. *see* Bruttomesso, D.
Pickkers, P. *see* Abbink, E.J.
Piovesan, A. *see* Procopio, M.
Platt, M.J., Stanisstreet, M., Casson, I.F., Howard, C.V., Walkinshaw, S., Pennycook, S. & McKendrick, O. St Vincent's Declaration 10 years on: outcomes of diabetic pregnancies, 216
van der Ploeg, H.M. *see* Adriaanse, M.C.
Polak, M. *see* Martin, D.
Poli, M. *see* Bonora, E.
van de Poll-Franse, L.V., Valk, G.D., Renders, C.M., Heine, R.J. & van Eijk, J.T.M. Longitudinal assessment of the developments of diabetic polyneuropathy and associated risk factors, 771
Porta, M. *see* Trento, M.
Poulsen, P. *see* Bentzen, J.
Poulton, J., Bednarz, A.L., Scott-Brown, M., Thompson, C., Macaulay, V.A., Simmons, D. The presence of a common mitochondrial DNA variant is associated with fasting insulin levels in Europeans in Auckland, 969

Pouwer, F. *see* DeVries, J.H.
Prasad, Y.D.M. *see* Oyibo, S.O.
Price, D.A., Lansang, M.C., Osei, S.Y., Fisher, N.D.L., Laffel, L.M.B. & Hollenberg, N.K. Type 2 diabetes, obesity, and the renal response to blocking the renin system with irbesartan, 858
Prins, A. *see* Frijling, B.D.
Pritchard, G.M. *see* Oyibo, S.O.
Prochazka, M. *see* Lindsay, R.S.
Procopio, M., Magro, G., Cesario, F., Piovesan, A., Pia, A., Molineri, N., Borretta, G. The oral glucose tolerance test reveals a high frequency of both impaired glucose tolerance and undiagnosed Type 2 diabetes mellitus in primary hyperparathyroidism, 958
Purevsuren, D. *see* Suvd, J.

Qabazard, M. *see* Shaltout, A.A.

Rachmani, R., Levi, Z., Slavachevski, I., Avin, M. & Ravid, M. Teaching patients to monitor their risk factors retards the progression of vascular complications in high-risk patients with Type 2 diabetes mellitus—a randomized prospective study, 385
Radder, J.K. *see* Broers, S.
Raffaelli, A. *see* Bonora, E.
Rajeswaran, C. ADA/WHO diabetes diagnostic criteria controversy—the Chennai chapter, 81
Ramachandran, A., Snehalatha, C., Vijay, V. & King, H. Impact of poverty on the prevalence of diabetes and its complications in urban southern India, 130
Ranstam, J. *see* Östgren, C.J.
Rao, S., Lindow, S.W. & Masson, E.A. Survey of pre-conception counselling, 615
Raoul, M. *see* Linkeschova, R.
Råstam, L. *see* Östgren, C.J.
Rave, K. *see* Heise, T.
Ravid, M. *see* Rachmani, R.
Raymond, N.T. *see* Hayward, L.M.
Rayner, C.K. *see* Horowitz, M
Reach, G. *see* Taverna, M.J.
Redekop, W.K., Koopmanschap, M.A., Rutten, G.E.H.M., Wolffenbuttel, B.H.R., Stolk, R.P. & Niessen, L.W. Resource consumption and costs in Dutch patients with Type 2 diabetes mellitus. Results from 29 general practices, 246
Rema, M., Saravanan, G., Deepa, R. & Mohan, V. Familial clustering of diabetic retinopathy in South Indian Type 2 diabetic patients, 910
Renders, C.M. *see* van de Poll-Franse, L.V.
Repetti, E. *see* Bo, S.
Reynolds, R.M. *see* Deepak, D.
Reynolds, R.M. *see* Hennessy, A.-R.
Rich, S.S. *see* Weijnen, C.F.
Richards, C. *see* Swift, B.
Richard, J.-L., Sultan, A., Daures, J.-P., Vannereau, D. & Parer-Richard, C. Diagnosis of diabetes mellitus and intermediate glucose abnormalities in obese patients based on ADA (1997) and WHO (1985) criteria, 292
Richmond, W., *see* Valabhji, J.
Riihimaa, P.H., Suominen, K., Knip, M., Tapanainen, P. & Tolonen, U. Cardiovascular autonomic reactivity is decreased in adolescents with Type 1 diabetes, 932
Rissanen, T. *see* Laaksonen, D.E.
Ritvanen, A. *see* Vääräsmäki, M.
Robert, J.J. *see* Martin, D.
Robinson, A.C.J. *see* Wahid, S.T.
Roderick, P. *see* Davies, M.
Rodríguez, I. *see* Abad, F.
Rogers, M. *see* Ko, G.T.C.
Roglic, G. *see* Suvd, J.
Rønningen, K.S. *see* Stene, L.C.
Rosenfalck, A.M., Almdal, T., Hilsted, J. & Madsbad, S. Body composition in adults with Type 1 diabetes at onset and during the first year of insulin therapy, 417
Rosskamp, R. *see* Heise, T.
Rotchford, A.P., Rotchford, K.M., Machattie, T. & Gill, G.V. Assessing diabetic control—reliability of methods available in resource poor settings, 195
Rotchford, K.M. *see* Rotchford, A.P.
Rotstein, R. *see* Elishkevitz, K.
Russel, F.G.M. *see* Abbink, E.J.
Rutten,G.E.H.M. *see* Redekop, W.K.
Ryder, C.H. *see* Abbott, C.A.

Saeed, B.O. *see* Wahid, S.T.
Sagen, J.V., Njølstad, P.R. & Søvik, O. Reduced prevalence of late-diabetic complications in MODY3 with early diagnosis, 697
Saggiani, F. *see* Bonora, E.
Saib, S.Z. *see* Chase, H.P.
Saito, T. *see* Nagasaka, S.
Sakai, M. *see* Nakai, Y.
Sakamoto, N. *see* Takeuchi, N.
Salonen, J.T. *see* Laaksonen, D.E.
Sampson, M.J., Shepstone, L., Greenwood, R.H., Harvey, I., Humphries, J., Heyburn, P.J., Temple, R.C. & Dole, G. An integrated mobile foot and retinal screening programme for people with Type 2 diabetes managed in primary care, 74
Samson, M. *see* Horowitz, M
Sano, T. *see* Takeuchi, N.
Sararols, L. *see* Hernández, C.
Saravanan, G. *see* Rema, M.
Särnblad, S., Kroon, M. & Åman, J. The short insulin tolerance test lacks validity in adolescents with Type 1 diabetes, 51
Sato, Y., Komatsu, M., Katakura, M., Ohfusa, H., Yamada, S., Yamauchi, K., Hiramatsu, K., Ichikawa, K., Aizawa, T. & Hashizume, K. Diminution of early insulin response to glucose in subjects with normal but minimally elevated fasting plasma glucose. Evidence for early β-cell dysfunction, 566
Sato, Y., Ohfusa, H., Katakura, M., Komatsu, M., Yamada, S., Yamauchi, K., Ichikawa, K., Aizawa, T. & Hashizume, K. A problem with the diagnosis of diabetes mellitus based on fasting plasma glucose, 82
Sawicki, P.T. *see* Didjurgeit, U.
Scaldaferri, E. *see* Bruttomesso, D.
Schachter, M. *see* Valabhji, J.
Schafer, A.J. *see* Berger, D.
van Schie, C.H.M. *see* Paisley, A.N.
Schmidt, L. *see* Garratt, A.M.
Schmitz, N. *see* Didjurgeit, U.
Schneider, B. *see* Maca, T.H.
Scott, A. Lipids and Vascular Disease Current Issues, 704

Scott-Brown, M. *see* Poulton, J.
Segura, R.M. *see* Hernández, C.
Seino, Y. *see* Nakai, Y.
Selam, J.-L. *see* Taverna, M.J.
Sera, Y. *see* Matsumoto, K.
Sera, Y. *see* Takino, H.
Serrano-Rios, M., Melchionda, N. & Moreno-Carretero, E. Role of sibutramine in the treatment of obese Type 2 diabetic patients receiving sulphonylurea therapy, 119
Seubert, M. *see* DeVries, J.H.
Shaltout, A.A., Moussa, M.A.A., Qabazard, M., Abdella, N., Karvonen, M., Al-Khawari, M., Al-Arouj, M., Al-Nakhi, A., Tuomilehto, J. & El-Gammal, A. Further evidence for the rising incidence of childhood Type 1 diabetes in Kuwait, 522
Shapira, I. *see* Elishkevitz, K.
Shapo, L., Coker, R. & McKee, M. Tracking diabetes in Albania: a natural experiment on the impact of modernization on health, 87
Shaw, J. *see* Unwin, N.
Shepherd, M. *see* Owen, K.R.
Shepstone, L. *see* Sampson, M.J.
Shigeta, H. *see* Kitamura, A.
Shikata, K. *see* Usui, H.
Shin, D.H. *see* Choi, K.M.
Shin, J. *see* McFarlane, S.I.
Shipley, M.J. *see* Batty, G.D.
Silink, M. *see* Kao, Y.
Simmons, D. *see* Poulton, J.
Simó, R. *see* Hernández, C.
Simpson, H. *see* Kipps, S.
Singh, B.M. *see* Buch, H.N.
Sinthanayothin, C., Boyce, J.F., Williamson, T.H., Cook, H.L., Mensah, E., Lal, S. & Usher, D. Automated detection of diabetic retinopathy on digital fungus images, 105
Skinner, T.C. *see* Howells, L.
Skinner, T.C. *see* Snoek, F.J.
Slama, G. *see* Taverna, M.J.
Slavachevski, I. *see* Rachmani, R.
Slee, F. *see* Cundy, T.
Smit, A.J. *see* Meijer, J.W.G.
Smits, P. *see* Abbink, E.J.
Snehalatha, C. *see* Ramachandran, A.
Snoek, F.J. *see* Adriaanse, M.C.
Snoek, F.J. *see* DeVries, J.H.
Snoek, F.J. & Skinner, T.C. Psychological counselling in problematic diabetes: does it help? 265
So, W.Y., Chan, J.C.N., Yeung, V.T.F., Chow, C.C., Ko, G.T.C., Li, J.K.Y., Cockram, C.S. Sulphonylurea-induced hypoglycaemia in institutionalized elderly in Hong Kong, 966
Sobngwi, E., Vexiau, P., Levy, V., Lepage, V., Mauvais-Jarvis, F., Leblanc, H., Mbanya, J.C. & Gautier, J.F. Metabolic and immunogenetic prediction of long-term insulin remission in African patients with atypical diabetes, 832
Sola, A. *see* Taverna, M.J.
Sonderen, E.V. *see* Meijer, J.W.G.
Song, Y.D. *see* Ahn, C.W.
Soos, M. *see* Berger, D.
Søvik, O. *see* Sagen, J.V.
Spijker, A.J. *see* DeVries, J.H.
Spinhoven, Ph. *see* Broers, S.
Spraul, M. *see* Linkeschova, R.
Stanisstreet, M. *see* Platt, M.J.
Stegmayr, B. *see* Eliasson, M.
Steinke, D.T. *see* Donnan, P.T.
Stene, L.C., Rønningen, K.S., Magnus, P. & Joner, G. Does HLA genotype influence the relative risk of Type 1 diabetes conferred by dietary factors? 429
Stenhouse, E.A. *see* Zhao, H.X.
Stephenson, C.R. *see* Feltbower, R.G.
Stevens, L.K. *see* Kempler, P.
Stokholm, K.H. *see* Ihlemann, N.
Stolk, R.P. *see* Redekop, W.K.
Stolk, R.P. *see* van der Feen, C.
Strachan, M.W.J., Gough, K., McKnight, J.A. & Padfield, P.L. Ambulatory blood pressure monitoring: is it necessary for the routine assessment of hypertension in people with diabetes, 787
Stride, A. *see* Owen, K.R.
Stückenschneider, P. *see* Didjurgeit, U.
Stümpflen, A. *see* Maca, T.H.
Sturgess, R. *see* Chong, J.S.W.
Sugimoto, T. *see* Usui, H.
Sultan, A. *see* Richard, J.-L.
Sultan, J. *see* Wahid, S.T.
Sun, B. *see* Herz, M.
Suominen, K. *see* Riihimaa, P.H.
Sutherland, W.H.F. *see* Manning, P.J.
Suvd, J., Gerel, B., Otgooloi, H., Purevsuren, D., Zolzaya, H., Roglic, G. & King, H. Glucose intolerance and associated factors in Mongolia: results of a national survey, 502
Svartberg, J. *see* Ahrén, B.
Swift, B., Hawkins, P.N., Richards, C. & Gregory, R. Examination of insulin injection sites: an unexpected finding of localized amyloidosis, 881

Tabák, Á.Gy., Kerényi, Zs., Nagy, E., Bosnyák, Zs., Madarász, E. & Tamás, Gy. Height and gestational diabetes mellitus, 344
Tack, C.J. *see* Abbink, E.J.
Tai, T.Y. *see* Chuang, L.-M.
Takemura, Y. *see* Aso, Y.
Takeuchi, N., Kawamura, T., Kanai, A., Nakamura, N., Uno, T., Hara, T., Sano, T., Sakamoto, N., Hamada, Y., Nakamura, J. & Hotta, N. The effect of cigarette smoking on soluble adhesion molecules in middle-aged patients with Type 2 diabetes mellitus, 57
Takino, H., Yamasaki, H., Abiru, N., Sera, Y., Abe, T., Kawasaki, E., Yamaguchi, Y., Eguchi, K., Kanazawa, Y. & Nagataki, S. Antibodies to GAD in Japanese patients classified as Type 2 diabetes at diagnosis. High titre of GAD Ab is a predictive marker for early insulin treatment—report of west Japan (Kyushu, Yamaguchi, Osaka) study for GAD Ab(+) diabetes, 730
Tam, S.C.F. *see* Tan, K.C.B.
Tam, W.H. *see* Ko, G.T.C.
Tamás, Gy. *see* Kempler, P.
Tamás, Gy. *see* Tabák, Á.Gy.
Tan, K.C.B., Tso, A.W.K., Tam, S.C.F., Pang, R.W.C., Lam, K.S.L. Acute effect of orlistat on post-prandial lipaemia and free fatty acids in overweight patients with Type 2 diabetes mellitus, 944

Tanaka, T. *see* Kitamura, A.
Taniguchi, A. *see* Nakai, Y.
Tapanainen, P. *see* Riihimaa, P.H.
Targher, G. *see* Bonora, E.
Taverna, M.J., Sola, A., Guyot-Argenton, C., Pacher, N., Bruzzo, F., Chevalier, A., Slama, G., Reach, G. & Selam, J.-L. eNOS4 polymorphism of the endothelial nitric oxide synthase predicts risk for severe diabetic retinopathy, 240
Tayama, K. *see* Aso, Y.
Taylor, R. *see* Pandit, R.J.
Temple, R.C. *see* Sampson, M.J.
Tesfaye, S. *see* Kempler, P.
Thomason, M.J. *see* Colhoun, H.M.
Thompson, C. *see* Poulton, J.
Tiengo, A. *see* Bruttomesso, D.
Tiengo, A. *see* Iori, E.
Tin-U, A. *see* Hulme, S.A.
Tolonen, U. *see* Riihimaa, P.H.
Toorians, A. *see* Hadithi, M.
Torkington, R. *see* Abbott, C.A.
Tracy, R.P. *see* Festa, A.
Trautner, C., Haastert, B., Giani, G. & Berger, M. Amputations and diabetes: a case-control study, 35
Trento, M., Bajardi, M., Borgo, E., Passera, P., Maurino, M., Gibbins, R., Owens, D.R., Cavallo, F. & Porta, M. Perceptions of diabetic retinopathy and screening procedures among diabetic people, 810
Tsai, S.T. *see* Chuang, L.-M.
Tso, A.W.K. *see* Tan, K.C.B.
Tubiana-Rufi, N. *see* Martin, D.
Tumini, S. *see* S. Mohn, A.
Tuomilehto, J. *see* Shaltout, A.A.
Tuvemo, T. *see* Tylleskär, K.
Tylleskär, K., Tuvemo, T. & Gustafsson, J. Author's reply, 700

Ueki, Y. *see* Matsumoto, K.
Ugurluoglu, A. *see* Maca, T.H.
Uno, T. *see* Takeuchi, N.
Unwin, N., Shaw, J., Zimmet, P. & Alberti, G. International Diabetes Federation IGT/IFG Consensus Statement. Report of an Expert Consensus Workshop 1–4 August 2001, Stoke Poges, UK, 708
Usher, D. *see* Sinthanayothin, C.
Usui, H., Shikata, K., Wada, J., Sugimoto, T., Yamana, J., Oishi, T., Matsuda, M., Yoneda, M., Koshima, I. & Makino, H. A case of congenital generalized lipodystrophy with lipoatrophic diabetes developing anti-insulin antibodies, 794

Vaag, A. *see* Bentzen, J.
Vääräsmäki, M., Gissler, M., Ritvanen, A. & Hartikainen, A.-L. Congenital anomalies and first life year surveillance in Type 1 diabetic births, 589
Valabhji, J., Donovan, J., McColl, A.J., Schachter, M., Richmond, W., & Elkeles, R.S. Rates of cholesterol esterification and esterified cholesterol net mass transfer between high-density lipoproteins and apolipoprotein B-containing lipoproteins in Type 1 diabetes, 424
Valbusa, D. *see* Iori, E.
Valk, G.D. *see* van de Poll-Franse, L.V.
Van Leberghe, S. *see* Hermans, M.P.
Van Ross, E.R.E. *see* Abbott, C.A.
Vangsness, L. *see* Anderson, B.J.
Vannereau, D. *see* Richard, J.-L.
van der Ven, N.C.W. *see* Broers, S.
Vexiau, P. *see* Sobngwi, E.
Vijay, V. *see* Ramachandran, A.
Visboll, T. *see* Koning, E.J.P.
van Vliet, K.P. *see* Broers, S.
Vora, J. Type 2 Diabetes in Practice, 171
Vora, J.P. *see* Younis, N.
Vuitch, F. *see* Haque, W.A.

Wada, J. *see* Usui, H.
Wahid, S.T., Sultan, J., Handley, G., Saeed, B.O., Weaver, J.U. & Robinson, A.C.J. Serum fructosamine as a marker of 5-year risk of developing diabetes mellitus in patients exhibiting stress hyperglycaemia, 543
Wakefield, A.J. *see* Montgomery, S.M.
Walker, J.D. & Warren, R.E. Necrobiosis lipoidica in Prader–Willi-associated diabetes mellitus, 884
Walker, J.D. *see* Deepak, D.
Walker, J.D. *see* Hennessy, A.-R.
Walker, J.D. *see* Hennessy, A.R.
Walkinshaw, S. *see* Platt, M.J.
Wallace, T.M. & Matthews, D.R. Coefficient of failure: a methodology for examining longitudinal β-cell function in Type 2 diabetes, 465
Wallace, T.M. & Matthews, D.R. The assessment of insulin resistance in man, 527
Wallymahmed, M. *see* Li Voon Chong, J.S.W.
Walton, J. *see* Kenardy, J.
Ward, J.D. *see* Kempler, P.
Warram, J.H. *see* Weijnen, C.F.
Warren, R.E. *see* Walker, J.D.
Warter-Neuhann, C. *see* Neuhann, H.F.
Wat, N.M.S. *see* Liu, Y.F.
Watkins, P. *see* Gill, G.
Watkins, P.J. *see* Bosman, D.R.
Weaver, J.U. *see* Wahid, S.T.
Webb, D.J. *see* Kempler, P.
Wedel, H. *see* Bokemark, L.
Weih, M.J. *see* Kuo, C.S.
Weijnen, C.F., Rich, S.S., Meigs, J.B., Krolewski, A.S. & Warram, J.H. Risk of diabetes in siblings of index cases with Type 2 diabetes: implications for genetic studies, 41
West, I.C. Glucose-6-phosphate dehydrogenase: a candidate gene for diabetes, 171
Whalley, A. *see* Chaturvedi, N.
Whalley, A.M. *see* Abbott, C.A.
Whitehead, J.P. *see* Berger, D.
Widdows, P. *see* Abbott, C.A.
Widdows, P. *see* Chaturvedi, N.
Wikstrand, J. *see* Bokemark, L.
Wilding, J.P.H. Neuropeptides and appetite control, 619
Wilkie, M. Diabetic Nephropathy, 704
Williams, R. *see* McFarlane, S.I.
Williams, R. *see* Winocour, P.H.
Williamson, S. *see* Abbott, C.A.
Williamson, T.H. *see* Sinthanayothin, C.
Wilson, A.C. *see* Howells, L.

Winocour, P.H., Ainsworth, A. & Williams, R. Association of British Clinical Diabetologists (ABCD) survey of secondary care services for diabetes in the UK, 2000. 1. Methods and major findings, 327
Wolffenbuttel, B.H.R. *see* Redekop, W.K.
van der Wouden, J.C. *see* Frijling, B.D.

Yamada, S. *see* Sato, Y.
Yamaguchi, M. *see* Kitamura, A.
Yamaguchi, Y. *see* Takino, H.
Yamana, J. *see* Usui, H.
Yamasaki, H. *see* Takino, H.
Yamauchi, K. *see* Sato, Y.
Yano, M. *see* Kamiuchi, K.
Yano, M. *see* Kitamura, A.
Yates, A.P. & Laing, I. Age-related increase in haemoglobin A_{1c} and fasting plasma glucose is accompanied by a decrease in β cell function without change in insulin sensitivity: evidence from a cross-sectional study of hospital personnel, 254
Yeo, G. *see* Berger, D.
Yeung, V.T.F. *see* So, W.Y.
Yomtov, B. *see* Perlemuter, L.
Yoneda, M. *see* Usui, H.
Yoshikawa, T. *see* Kamiuchi, K.
Yoshikawa, T. *see* Kitamura, A.
Young, R.J. *see* Bhattacharyya, A.
Younis, N., Broadbent, D.M., Harding, S.P., Vora, J.P. Prevalence of diabetic eye disease in patients entering a systematic primary care-based eye screening programme, 1014
Yung, B., Noormohamed, F.H., Kemp, M., Hooper, J., Lant, A.F. & Hodson, M.E. Cystic fibrosis-related diabetes: the role of peripheral insulin resistance and β-cell dysfunction, 221
Zasloff, M. *see* Ge, Y.
Zeltser, D. *see* Elishkevitz, K.
Zenari, L. *see* Bonora, E.
Zenere, M.B. *see* Bonora, E.
Zhao, H.X., Moyeed, R.A., Stenhouse, E.A., Demaine, A.G. & Millward, B.A. Space–time clustering of childhood Type 1 diabetes in Devon and Cornwall, England, 667
Zimmet, P. *see* Unwin, N.
Zolzaya, H. *see* Suvd, J.

Subject Index

16189 variant, 681

accumulation, 490
ADA, 292
ADA fasting criteria, 853
adherence, 279, 635, 643, 685
adhesion molecule, 371, 822
adhesion molecules, 57
adolescence, 643, 649, 932
adolescents, 212, 827
adult, 611
adult-onset, 986
aetiology, 758
Africa, 195, 895
African, 832
Africans, 575
afternoon meal, 1026
age at onset of diabetes, 924
age, 254, 558
age-adjustment, 954
albumin excretion rate, 901
aldose reductase, 113
ambulatory blood pressure monitoring, 787
amputation, 99, 575
amputations, 35
anaemia, 65
angiography, 366
ankle pressure, 307
anti-oxidant status, 752
antibodies, 518
Apolipoprotein B, 12
arthroplasty, 338
Asia population, 978
Asia, 549
assessment, 527
atherosclerosis, 144, 889, 939
atorvastatin, 201
attractor, 274
attributable risk, 35
atypical diabetes, 832
audit with feedback, 836
autonomic nervous system, 932
autonomic neuropathy, 177

β cell, 254, 465
β-cell dysfunction, 566
β-cell function, 221
bifurcation, 274
blindness, 27
Blood Glucose Awareness Training, 157
blood glucose monitoring, 635
blood pressure, 359
body composition, 417
body mass index, 558, 681
bolus insulin, 317

C-peptide, 227
C-reactive protein, 689, 847, 939
cardiovascular disease, 901
cardiovascular reflex tests, 932
cardiovascular risk, 549
CARDS, 201
care, 509
carotid atherosclerosis, 57
case-control study, 35
catecholamine, 359
Caucasians, 41
cell adhesion molecules, 847
central fat, 689
central obesity, 502
chaos, 274
chaotic control, 274
Charcot neuroarthropathy, 338
CHD, 580
chest pain, 862
childhood, 70
chidhood diabetes, 1026
childhood diabetes, 162
childhood Type 1 diabetes, 667
children, 518, 827
Chinese, 113
cholesterol, 514
cholesterol esterification, 424
chronic hypertension, 482
chronic venous ulcer, 440
cigarette smoking, 57
circadian rhythm, 359
clamp, 527
clinical decision making, 836
clinical presentation, 509
clinical trial, 234
clot retraction, 862
coefficient, 465
coeliac disease, 334
comparison, 661
complications, 35, 246, 814, 843
complications of diabetes, 130
computer simulation, 762
congenital anomalies, 322
congenital anomaly, 589
congenital malformations, 216

continuous intraperitoneal insulin infusion, 496
continuous subcutaneous insulin infusion, 628
coronary artery disease, 862
coronary heart disease, 424, 470
coronary heart disease risk, 514
costs, 246, 594
counselling, 265
CSII, 746
cystic fibrosis, 221

deafness, 89
developing countries, 895
diabetes, 27, 35, 130, 195, 201, 221, 509, 514, 518, 549, 558, 572, 575, 611, 787, 810, 814, 836, 847, 853, 858, 867, 889, 895, 954
diabetes education, 799
diabetes management, 978
diabetes mellitus, 1, 152, 167, 265, 292, 307, 338, 366, 385, 412, 435, 440, 456, 502, 689, 735, 762, 784, 862, 874, 958, 1014
diabetes mellitus type 2, 300, 949
diabetes professionals, 799
diabetes services, 327
diabetes specialist nurses, 327
diabetes teamwork, 799
diabetic autonomic neuropathy, 65, 901
diabetic complications, 385
diabetic diagnosis, 105
diabetic foot, 400
diabetic foot care, 575
diabetic foot ulcer, 440
diabetic nephropathy, 1000
diabetic neuropathy, 19, 440, 661, 827, 962
diabetic polyneuropathy, 771
diabetic pregnancy, 589
diabetic retinopathy, 240, 285, 371, 655, 762, 810, 910, 924, 1014
diagnosis, 177, 406, 843
diagnostic criteria, 292, 549
dieticians, 327
DNE score, 962
DNS score, 962
driving, 435
driving and hypoglycaemia, 157
DXA scanning, 417
dyslipidaemia, 535

eating disorder, 234
effective learning in diabetes, 799
elderly, 949
employment, 435
endothelial dysfunction, 19
endothelial function, 476
energy balance, 619
eNOS4 polymorphism, 240
epidemiology, 35, 167, 502, 522, 973
erythrocyte adhesiveness/aggregation, 152
erythropoietin, 65
esterified cholesterol net mass transfer, 424
ethnic diferences, 973
ethnicity, 162
euglycaemic clamp, 51
Europe, 594
evalence, 954

failure, 465
familial clustering, 910
familial, 1022
families, 994
family conflict, 635
family history, 689
family study, 735
fasting glucose, 681
fixed insulin lispro mixtures, 918
foot, 1009
foot infection, 895
foot ulcer, 377
foot ulceration, 19
forced expiratory volume, 167
Framingham equation, 514
free fatty acids, 944

GAD Ab, 730
GADA, 227
gastric emptying, 177
gastrointestinal dysmotility, 611
gastrostomy, 1006
gene, 113
general practice, 836
genetic, 240
genetics, 89, 994
geographical distribution, 522
gestational diabetes, 322, 351
gestational hypertension, 482
glibenclamide, 136, 673
gliclazide, 752
glimepiride, 136
glucose, 195, 1006
  analysers, 693
  excursions, 870
  homeostasis, 254
  metabolism, 535
  tolerance test, 351, 735
  toxicity, 125
glutamic acid decarboxylase, 602
glyburide combination therapy, 673
glycaemic control, 157, 177, 274, 393, 635, 924, 978
glycosylated haemoglobin, 195

hand infection, 895
hand sepsis, 895
HbA_{1c}, 465, 978
health perceptions, 810
health planning, 762
heart rate variability, 359, 932
heritability, 994
heteroplasmy, 784
high-density lipoproteins, 424
HLA, 832
HNF-1α, 758
HOMA, 221, 470
homeostatic model assessment, 527
hormone replacement therapy, 847
hyperglycaemia, 611
hypertension, 502, 535, 787
hypoglycaemia, 435, 746, 918
hypoglycaemic awareness, 157
hypothalamus, 619
hypoxia, 65

ICAM-1, 371
Icodextrin metabolites, 693
IL-6 gene polymorphism, 1000
image analysis, 105
image recognition, 105
immunization, 986
impaired fasting glycaemia, 292, 456
impaired glucose regulation, 558
impaired glucose tolerance, 292, 502, 958
in-patient management, 412
incidence, 162, 377, 522
India, 130
infant mortality, 589
infection, 986
inflammation, 939
iNOS, 777
insulin, 12, 144, 572, 724, 777, 1006
  analog, 1026
  analogue, 393, 918
  aspart, 393
  glargine, 490
  infusion systems, 496
  pump therapy, 746
  receptor substrate-1, 804
  remission, 832
  resistance, 221, 470, 527, 535, 689, 724, 804, 939, 958, 994
  resistance syndrome, 724
  resistence, 973
  secretion, 832
  sensitivity, 51, 254, 527, 735
  therapy, 490
  treatment, 730
intensified insulin therapy, 814
intensive management, 385
interleukin-6, 1000
intima-media thickness, 144
intussusception, 611
islet amyloidosis, 1022

Japanese, 566

K-function, 667
K_{ATP} channels, 136

ketoacidosis, 777
knee, 338
Knox's test, 667
Korean, 227, 681, 853

lamin A/C, 1022
leptin, 619
light plethysmography, 307
limb sepsis, 895
limited joint mobility, 1009
linoleate, 456
lipase inhibitor, 944
lipid-lowering, 201
lipodystrophy, 1022
lipoprotein(a), 300
long-acting insulin analogue, 490
lower extremity atherosclerosis, 366
lymphomonocyte, 777

macroalbuminuria, 1000
macroangiopathy, 822
markers for diabetes risk, 543
mass screening, 1014
maternal inheritance, 89
men, 580
metabolic control, 70
metabolic syndrome, 724
metformin, 136, 673
MI, 448
microalbuminuria, 212, 1000
microangiopathy, 822
microvascular complications, 973
mitochondrial DNA, 89, 681
mitochondrial gene mutation, 784
model, 527
MODY, 758
Mongolia, 502
MONICA, 874
mortality, 448, 470, 580
multiple daily injections, 628
multiprofessional education, 799
multiprofessional learning, 799

nephropathy, 65, 113
neural network, 105
neurological examination, 771
neuropathy disability score, 377
neuropathy, 400
neuropeptide, 619
neuropsychological function, 311
nitric oxide, 240, 655, 777
non-linear dynamics, 274
nutrition, 417

obese, 572
obesity, 41, 119, 292, 535, 619, 858
odds ratio, 35
older patients, 311
Oman, 954
ophthalmoscopy, 867
optometrists, 741
oral anti-diabetic therapy, 673
oral glucose tolerance test, 292, 566, 853
oral hypoglycaemic agents, 279, 606, 685
outreach visits, 836

pain, 827
painful neuropathy, 870
paraoxonase, 212
parental involvement, 635
patient-assessed, 1
patient education, 385
peak pressure, 1009
perceptions, 406
percutaneous transluminal angioplasty, 300
perinatal mortality, 216, 589
peripheral arterial occlusive disease, 300
peripheral neuropathy, 74, 575
peripheral vascular disease, 74, 99
pertussis, 986
pharmacokinetics, 490
photography, 1014
physical activity, 580
physicians, 327
plantar aponeurosis, 1009
plasma lipids, 901
platelet function, 862
podiatrists, 327
polycystic ovaries, 1022
polymorphism, 12, 113, 371
polyneuropathy, 962
polyunsaturated fatty acids, 456
population, 279, 377, 448
population-based, 27
population survey, 874
post-challenge plasma glucose, 939
post-prandial glucose, 317
poverty, 130
preeclampsia, 482
pregnancy, 216, 482
premixed insulin, 393
pressure time integrals, 1009
prevalence, 558, 874, 954, 1014
prevention, 594
primary care, 74, 125
primary health care, 771
primary hyperparathyroidism, 958
pro-opiomelanocortin, 619
problem solving, 643
procedure, 406
prospective study, 771
pseudo-hyperglycaemia, 693
psychology, 234, 265
psychological and psychosocial problems, 814
psychotherapy, 814
puberty, 51, 932
pulseoximetry, 307

quality, 594
quality assurance, 285
quality of life, 1, 496, 628, 746
quantitative sensory testing, 400
quantitative sensory threshold, 661

random serum glucose, 543
registers, 27
relative risk, 35
reliability, 1
renal failure, 889
renal function, 858
renin, 858
renovascular disease, 889
reproducibility, 514
restenosis, 300
retinal photography, 867
retinal screening, 867
retinopathy, 105, 212, 602, 741
review, 1
risk factor, 377
risk factors, 99
rosiglitazone, 572

satisfaction survey, 412
saturated fatty acids, 456
screen, 377, 400
screening, 74, 285, 292, 351, 406, 518, 543, 549, 741, 762, 810
self efficacy, 643
self-care behaviour, 157
sensory thresholds, 827
serum fructosamine, 543
serum insulin, 125
short insulin tolerance test, 51
sibling recurrence risk, 41
sibutramine, 119
skin, 1009
slit-lamp biomicroscopy, 741
SNRI, 119
socio-economic status, 843
socioeconomic factors, 130
South Asian, 99
South Indians, 910
South-west of England, 667
space–time clustering, 667
stomach, 177
stress hyperglycaemia, 543
stroke, 1006
subclinical hypothyroidism, 70
surveillance, 522
symptoms, 177
symptom score, 962
syndromes, 804

Tanzania, 509
therapy, 177
thiazolidinedione, 572
thresholds, 465
thyroid, 334
thyroiditis, 518
time to insulin, 606

time trend, 874
time trends, 162
TNF-α, 949
toe pressure, 307
total body water, 417
transforming growth factor, 440
transition arrangements, 649
treatment options, 509
trends, 954
trial, 201
triglyceride, 12, 944
triglyceriderich lipoprotein remnants, 944
tropical diabetic hand syndrome, 895
twice-daily regimen, 393
Type 1 (insulin-dependent) diabetes mellitus, 901
Type 1 diabetes, 51, 70, 212, 216, 227, 317, 322, 334, 417, 424, 490, 522, 628, 635, 643, 777, 918, 1026
Type 1 diabetes mellitus, 924, 986
Type 2 diabetes, 12, 41, 113, 119, 227, 234, 246, 279, 311, 322, 406, 448, 476, 482, 535, 580, 606, 673, 685, 730, 771, 843, 910
Type 2 diabetes mellitus, 57, 74, 594, 602, 752, 822, 944
Type 2 diabetic patient, 359

ultrasound, 518, 1009

validity, 1
variants, 804
vascular complications, 476
vascular endothelial growth factor, 655
vascular function, 752
vasodilatation, 777
venous occlusion plethysmography, 136
vertical transmission, 41
vibration perception threshold, 661
visual impairment, 27
vital capacity, 167
vitreous fluid, 655
von Willebrand factor, 19, 476

WHO, 292
WHO 2-h criteria, 853

young age onset diabetes, 227
young-onset Type 2 diabetes, 758
YT2D, 758